

RESULTS OF THE RADIOLOGICAL SURVEY AT 79 AVENUE B, LODI,

062798

ORNL/RASA-88/79

M-055

NEW JERSEY (LJ091)

> R. D. Foley L. M. Floyd

OPERATED BY MARTIN MARIETTA ENERGY SYSTEMS, INC. FOR THE UNITED STATES DEPARTMENT OF ENERGY This report has been reproduced directly from the best available copy.

•

Available to DOE and DOE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831; prices evailable from (615) 576-8401, FTS 626-8401.

8062798

1.00

Available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22181. NTIS price codes-Printed Copy: A03 Microfiche A01

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

ORNL/RASA-88/79

ことをいいていたい

HEALTH AND SAFETY RESEARCH DIVISION

s

Nuclear and Chemical Waste Programs (Activity No. AH 10 05 00 0; ONLWCO1)

RESULTS OF THE RADIOLOGICAL SURVEY AT 79 AVENUE B, LODI, NEW JERSEY (LJ091)

R. D. Foley and L. M. Floyd

Date Published - June 1989

Investigation Team

R. E. Swaja – Measurement Applications and Development Manager W. D. Cottrell – FUSRAP Project Director R.D. Foley – Field Survey Supervisor

Survey Team Members

A. C. Butler W. H. Shinpaugh^{*} P. F. Tiner W. Winton *Stone Associates

Work performed by the MEASUREMENT APPLICATIONS AND DEVELOPMENT GROUP

Prepared by the OAK RIDGE NATIONAL LABORATORY Oak Ridge, Tennessee 37831-6285 operated by MARTIN MARIETTA ENERGY SYSTEMS, INC. for the U. S. DEPARTMENT OF ENERGY under Contract No. DE-AC05-84OR21400

al - Sala a state

CONTENTS

4,

062798

LIST OF FIGU	RES .	• •	•	•	٠	٠	•	٠	٠	•	•	•	•	٠	٠	٠	•	•	٠	٠	٠	•	•	٠	v
LIST OF TABI	LES .	• •	٠	•	•	•	•	•	٠	•	•	•	٠	•	٠	•	•	٠	•	•	•	•	•	•	vii
ACKNOWLED	GMEN	TS	•	•	•	•	•	٠	•	•	•	٠	•	•	•	•	٠	•	٠	•	•	•	•	•	ix
ABSTRACT .	• • •	• •	•	•	•	٠	٠	٠	•	•	•	٠	•	٠	•	٠	•	•	•	•	•	•	•	•	xi
INTRODUCTI	ON .	.••	•	•	٠	•	•	٠	•	•	٠	٠	•	•	•	•	•	•	•	•	•	•	•	•	1
SURVEY MET	HODS		•	•	•	•	•	٠	•	•	•	•	•	•	•	•	•	•	٠	•	•	•	٠	•	2
SURVEY RESU	ULTS.	••	•	•	•	•	٠	•	•	•	•	•	•	•	•	٠	•	•	•	٠	•	•	•	•	2
Gamma Ra	diation	Lev	els	•	•	•	٠	٠	•	•	٠	•	•	٠	•	•	•	•	•	•	•	•	•	•	2
Systematic	and Bi	ased	So	il	Sa	m	ple	3	•	٠	٠	٠	•	٠	٠	•	•	٠	٠	•	•	•	•	٠	3
SIGNIFICANC	EOF	FINI	DIN	G	S	•	٠	•	•	•	٠	•	•	•	•	•	•	•	•	•	•	•	•	•	3
REFERENCES	S					•	•	•	•	•		•	•	•	•	•							•	•	4

LIST OF FIGURES

and a strate of the second second second

. مىدىرە

in the second

1

1

.

 $\left|\cdot\right\rangle$

1	Gamma radiation levels $(\mu R/h)$ measured on the surface at 79 Avenue B, Lodi, New Jersey (LJ091), with corresponding measurements one meter above the surface where indicated.	5
2	Diagram showing locations of soil samples taken at 79 Avenue B, Lodi, New Jersey (LJ091)	6

LIST OF TABLES

052793

••

ころうれたいとう いっていましょう 大学 まとうなま たままままたあっしょう

i,

÷

1	Applicable guidelines for protection against radiation
2	Background radiation levels for the northern New Jersey area
3	Concentrations of radionuclides in soil at 79 Avenue B, Lodi, New Jersey (LJ091)

vii

ACKNOWLEDGMENTS

Research for this project was sponsored by the Division of Facility and Site Decommissioning Projects, U.S. Department of Energy, under Contract No. DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc. The authors wish to acknowledge the support of J. E. Baublitz, Acting Director, Office of Remedial Action and Waste Technology; J. J. Fiore, Director, Division of Facility and Site Decommissioning Projects; and members of their staffs. The authors also appreciate the contributions of J. L. Rich, S. W. Hawthorne, B. C. Littleton, and L. J. Jeffers of the Publications Division; M. S. Uziel of the Environmental Information Systems Group; B. S. Ellis, D. A. Roberts, and T. R. Stewart of the Measurement Applications and Development Group; and W. H. Shinpaugh of Don Stone Associates for participation in the collection, analyses, editing, and reporting of data for this survey.

ABSTRACT

Maywood Chemical Works (MCW) of Maywood, New Jersey, generated process wastes and residues associated with the production and refining of thorium and thorium compounds from monazite ores from 1916 to 1956. MCW supplied rare earth metals and thorium compounds to the Atomic Energy Commission and various other government agencies from the late 1940s to the mid-1950s. Area residents used the sandlike waste from this thorium extraction process mixed with tea and cocoa leaves as mulch in their yards. Some of these contaminated wastes were also eroded from the site into Lodi Brook. At the request of the U.S. Department of Energy (DOE), a group from Oak Ridge National Laboratory conducts investigative radiological surveys of properties in the vicinity of MCW to determine whether a property is contaminated with radioactive residues, principally ²²²Th, derived from the MCW site. The survey typically includes direct measurement of gamma radiation levels and soil sampling for radionuclide analyses. The survey of this site, 79 Avenue B, Lodi, New Jersey (LJ091), was conducted during 1988.

Results of the survey demonstrated radionuclide concentrations in excess of the DOE Formerly Utilized Sites Remedial Action Program criteria. The radionuclide distributions are typical of the type of material originating from the MCW site.

RESULTS OF THE RAPIOLOGICAL SURVEY AT 79 AVENUE B, LODI, NEW JERSEY (LJ091)*

INTRODUCTION

From 1916 to 1956, process wastes and residues associated with the production and refining of thorium and thorium compounds from monazite ores were generated by the Maywood Chemical Works (MCW), Maywood, New Jersey. During the latter part of this period, MCW supplied rare earth metals and thorium compounds to various government agencies. In the 1940s and 1950s, MCW produced thorium and lithium, under contract, for the Atomic Energy Commission (AEC). These activities ceased in 1956, and, approximately three years later, the 30-acre real estate was purchased by the Stepan Company. The property is located at 100 Hunter Avenue in a highly developed area in Maywood and Rochelle Park, Bergen County, New Jersey.

During the early years of operation, MCW stored wastes and residues in lowlying areas west of the processing facilities. In the early 1930s, these areas were separated from the rest of the property by the construction of New Jersey State Highway 17. The Stepan property, the interim storage facility, and several vicinity properties have been designated for remedial action by the U.S. Department of Energy (DOE).

The waste produced by the thorium extraction process was a sandlike material containing residual amounts of thorium and its decay products, with smaller quantities of uranium and its decay products. During the years 1928 and 1944 to 1946, area residents used these process wastes mixed with tea and cocoa leaves as mulch in their lawns and gardens. In addition, some of the contaminated wastes were apparently eroded from the site into Lodi Brook and carried downstream.

Lodi Brook is a small stream flowing south from Maywood with its headwaters near the Stepan waste storage site. Approximately 150 ft after passing under State Route 17, the stream has been diverted underground through concrete or steel culverts until it merges with the Saddle River in Lodi, New Jersey. Only a small section near Interstate 80 remains uncovered. From the 1940s to the 1970s when the stream was being diverted underground, its course was altered several times. Some of these changes resulted in the movement of contaminated soil to the surface of a few properties, where it is still in evidence. In other instances, the contaminated soil was covered over or mixed with clean fill, leaving no immediate evidence on the surface. Therefore, properties in question may be drilled in search of former stream bed material, even in the absence of surface contamination.

As a result of the Energy and Water Appropriations Act of Fiscal Year 1984, the property discussed in this report and properties in its vicinity contaminated with residues from the former MCW were included as a decontamination research

^{*}The survey was performed by members of the Measurement Applications and Development Group of the Health and Safety Research Division at Oak Ridge National Laboratory under DCE contract DE-AC05-84OR21400.

and development project under the DOE Formerly Utilized Sites Remedial Action Program. As part of this project, DOE is conducting radiological surveys in the vicinity of the site to identify properties contaminated with residues derived from the MCW. The principal radionuclide of concern is thorium-232. The radiological surveys discussed in this report are part of that effort and were conducted, at the request of DOE, by members of the Measurement Applications and Development Group of the Oak Ridge National Laboratory.

2

A radiological survey of the private, residential property at 79 Avenue B, Lodi, New Jersey, was conducted during 1988. The survey and sampling of the ground surface were carried out on May 10, 1988.

SURVEY METHODS

The radiological survey of the property included: (1) a gamma scan of the entire property outdoors, both at the surface and one meter above the surface, and (2) collection of soil samples. No indoor survey measurements were performed.

Using a portable gamma scintillation meter, ranges of measurements were recorded for areas of the property surface and one meter above the surface. If the gamma readings were elevated, a biased soil sample was taken at the point showing the highest gamma radiation level. Systematic soil samples were taken at various locations on the property, irrespective of gamma radiation levels. These survey methods followed the plan outlined in Reference 1. A comprehensive description of the survey methods and instrumentation has been presented in another report.²

SURVEY RESULTS

Applicable federal guidelines are summarized in Table 1.³ The normal background radiation levels for the northern New Jersey area are presented in Table 2. These data are provided for comparison with survey results presented in this section. All direct measurement results presented in this report are gross readings; background radiation levels have not been subtracted. Similarly, background concentrations have not been subtracted from radionuclide concentrations measured in environmental samples.

Gamma Radiation Levels

Gamma radiation levels measured during a gamma scan of the surface of the property are given in Fig. 1, with corresponding measurements one meter above the surface where indicated. Gamma exposure rates over the major portion of the property surface ranged from 4 to 9 μ R/h. One-meter readings were between 6 and 26 μ R/h, with the 11- and 26- μ R/h readings being above the normal background level of 8 μ R/h at one meter for the northern New Jersey area (Table 2). The highest readings were in the eastern corner of the backyard, ranging from 11 to 85 μ R/h at the surface, and from the foundation of the house on the south side, reading 11 μ R/h.

Systematic and Biased Soil Samples

Systematic and biased soil samples were taken from various locations on the property for radionuclide analyses. Locations of the systematic (S) and biased (B) samples are shown in Fig. 2, with results of laboratory analyses provided in Table 3. Concentrations of radium, thorium, and uranium in these samples ranged from 0.61 to 5.1 pCi/g, 0.69 to 64 pCi/g, and 0.49 to <2.5 pCi/g, respectively. Biased samples B1A and B1B were above DOE guidelines for 232 Th (Table 1).

SIGNIFICANCE OF FINDINGS

Measurements taken at 79 Avenue B indicate that the property contained radioactive contamination primarily from the 232 Th and 238 U decay chains, with slight contamination from 226 Ra. These radionuclide distributions are typical of the type of material originating from the MCW site. The concentration and extent of 232 Th on this property were in excess of the applicable DOE criteria (Table 1). As shown in Fig. 2, this material was found at sample location B1, consistent with the elevated gamma radiation measurements in that area. Based on the results of this radiological assessment, it is recommended that this site be considered for inclusion in the DOE remedial action program.

REFERENCES

1. T. S

10

144

- 1. W. D. Cottrell, ORNL, to A. J. Whitman, DOE/HQ, correspondence, "Radiological Survey of Private Properties in Lodi, New Jersey" (August 15, 1984).
- 2. T. E. Myrick, B. A. Berven, W. D. Cottrell, W. A. Goldsmith, and F. F. Haywood, Procedures Manual for the ORNL Radiological Survey Activities (RASA) Program, Oak Ridge National Laboratory, ORNL/TM-8600 (April 1987).
- 3. U.S. Department of Energy, Guidelines for Residual Radioactivity at Formerly Utilized Sites, Remedial Action Program and Remote Surplus Facilities Management Program Sites (Rev. 2, March 1987).
- 4. U.S. Department of Energy, Radiological Survey of the Middlesex Municipal Landfill, Middlesex, New Jersey, DOE/EV-00005/20(April 1980).
- 5. T. E. Myrick and B. A. Berven, State Background Radiation Levels: Results of Measurements Taken During 1975-1979, Oak Ridge National Laboratory, ORNL/TM-7343 (November 1981).



79 Avenue B

Fig. 1. Gamma radiation levels $(\mu R/h)$ measured on the surface at 79 Avenue B, Lodi, New Jersey (LJ091), with corresponding measurements one meter above the surface where indicated.



Ę



Table 1.	Applicable	guidelines for	protecti	on agai	inst rad	liation*
----------	------------	----------------	----------	---------	----------	----------

062

798

Mode of exposure	Exposure conditions	Guideline value
Radionuclide concen- trations in soil	Maximum permissible con- centration of the follow- ing radionuclides in soil above background levels averaged over 100 m ² area ²³² Th ²³⁰ Th ²³⁸ Ra ²²⁶ Ra	5 pCi/g averaged over the first 15 cm of soil below the surface; 15 pCi/g when averaged over 15-cm thick soil layers more than 15 cm below the surface

*Reference 3.

Table 2.	Background radia	tion levels for tl	1e
	northern New Jer	sey area	

Type of radiation measurement or sample	Radiation level or radionuclide concentration
Gamma exposure at 1 m above ground surface $(\mu R/h)$	8ª
Concentration of radionuclides in soil (pCi/g) ²³² Th ²³⁸ U ²²⁶ Ra	0.9 ⁸ 0.9 ⁸ 0.9 ⁸

P

*Reference 4. *Reference 5.

1.100

•		Radionuclide concentration (pCi/g)							
Sample ^e	Depth (cm)	226 Ra	²³² Th ^{\$}	238 U ^{\$}					
<u></u>		Systematic sam	ples ^c						
S1A S2A	0-15 0-15	0.61±0.01 0.66±0.02	0.85±0.02 0.69±0.03	<2.0 <2.5					
		Biased sample	54						
B1A B1B B1C B1D B2A B2B	0-15 15-30 30-45 45-60 0-15 15-30	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\begin{array}{rrrr} 64 & \pm 2 \\ 22 & \pm 0.8 \\ 8.3 & \pm 0.5 \\ 3.5 & \pm 0.07 \\ 4.4 & \pm 0.2 \\ 3.1 & \pm 0.2 \end{array}$	• <2.0 0.49±3 1.3 ±2 1.1 ±3 <1.6 ±3					

Table 3. Concentrations of radionuclides in soil at 79 Avenue B, Lodi, New Jersey (LJ091)

8

^eLocations of soil samples are shown on Fig. 2.

³Indicated counting error is at the 95% confidence level $(\pm 2\sigma)$. ^cSystematic samples are taken at locations irrespective of gamma expo-

Biased samples are taken from areas shown to have elevated gamma exposure rates.

*Sample was not analyzed for ²³⁸U.

ORNL/RASA-88/79

062798

INTERNAL DISTRIBUTION

B. A. Berven
R. F. Carrier
W. D. Cottrell
A. G. Croff
J. W. Crutcher
L. M. Floyd
7-11. R. D. Foley
S. V. Kaye

- 13. P. Y. Lu 14. P. T. Owen 15-17. R. E. Swaja 18. J. K. Williams
 - 19. Central Research Library 20. IR&A Publications Office
 - 21. Laboratory Records RC
 - 22. Y-12 Technical Library

EXTERNAL DISTRIBUTION

- 23. J. D. Berger, Oak Ridge Associated Universities, P.O. Box 117, Oak Ridge, TN 37831
- 24. R. W. Doane, Eberline, Inc., 800 Oak Ridge Turnpike, P.O. Box 350, Oak Ridge, Tn 37831
- 25. J. J. Fiore, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
- 26-28. G. K. Hovey, Bechtel National, Inc., 800 Oak Ridge Turnpike, P.O. Box 350, Oak Ridge, TN 37831
 - 29. L. R. Levis, Roy F. Weston, Inc., 20030 Century Blvd., Germantown, MD 20874
 - 30. G. P. Turi, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
 - 31. J. W. Wagoner, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
- 32-34. Andrew Wallo III, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
 - 35. Office of Assistant Manager, Energy Research and Development, Oak Ridge Operations Office, P.O. Box 2001, Oak Ridge, TN 37831-8600
- 36-37. Office of Scientific and Technical Information, DOE, P.O. Box 62, Oak Ridge, TN 37831